arXiv:1304.7633v1 [q-bio.BM] 29 Apr 2013

Molecular Dynamics Studies on the NMR Structures of Rabbit Prion Protein Wild-type and Mutants

Jiapu Zhang[1*], **Yuanli Zhang**[2]
[1]Graduate School of Sciences, Information Technology and Engineering, CIAO, The University of Ballarat, MT Helen Campus, Victoria 3353, Australia;
[2]School of Basic Medical Sciences, Taishan Medical University, Shandong 271000, China.
[*]Correspondence address:
Tel: +61-3-5327 6335, 0423 487 360; Email: j.zhang@ballarat.edu.au, jiapu_zhang@hotmail.com

Abstract:
Prion diseases, traditionally referred to as transmissible spongiform encephalopathies (TSEs), are invariably fatal and highly infectious neurodegenerative diseases that affect a wide variety of mammalian species, manifesting as scrapie in sheep and goats, bovine spongiform encephalopathy (or mad-cow disease) in cattle, chronic wasting disease in deer and elk, and Creutzfeldt-Jokob diseases, Gerstmann-Strussler-Scheinker syndrome, fatal familial insomnia, and kulu in humans, etc. These neurodegenerative diseases are caused by the conversion from a soluble normal cellular prion protein ($\mathrm{PrP^C}$) into insoluble abnormally folded infectious prions ($\mathrm{PrP^{Sc}}$) and the conversion of $\mathrm{PrP^C}$ to $\mathrm{PrP^{Sc}}$ is believed to involve conformational change from a predominantly α-helical protein to one rich in β-sheet structure. Such conformational changes may be amenable to study by molecular dynamics (MD) techniques. For rabbits, classical studies show they have a low susceptibility to be infected by $\mathrm{PrP^{Sc}}$, but recently it was reported that rabbit prion can be generated through PMCA in vitro and the rabbit prion is infectious and transmissible (Proc Natl Acad Sci USA 109(13) 5080-5085 (2012)). This paper studies the NMR structure of rabbit prion protein wild-type and mutants by MD techniques, in order to understand the specific mechanism of rabbit prion protein and rabbit prions.

Keywords: prion diseases; rabbit prion proteins; rabbit prions; NMR structures; molecular dynamics study

1 Introduction

“Mad cow disease” (Bovine Spongiform Encephalopathy (BSE)) belongs to a contagious type of Transmissible Spongiform Encephalopathy (TSE). Scientists believe it is caused by Prions (the misfolding prion proteins) but they may have not yet solved the riddle of “mad cow disease”. This is due to a Prion is neither a virus, a bacteria nor any microorganism so the disease cannot be caused by the vigilance of the organism immune system and it can freely spread from one species to another species. The humans exists the susceptibility of TSEs. For example, the human version of mad cow disease named

Creutzfeldt-Jakob Disease (CJD) and variant CJD (vCJD) just happen randomly through infections of transplanted tissue or blood transfusions or consumption of infected beef products. Cats, mink, deer, elk, sheep and many other animals are also susceptible to TSEs. However, rabbits, horses and dogs seem to be unaffected by Prions [16, 7, 13]. Scientists do not know the reason.

The prion protein is a naturally occurring protein in vivo. Its lesions in brain are not caused by the vigilance of the immune system. Recent studies have found that the lesions led astray as long as by contact with other normal prion proteins. The cells are arranged in accordance with the instruction of the gene and formed into proteins with different shapes and functions. But, like cardboard boxes, proteins need to be properly "folded" in order to ensure their normal work. When proteins are folded into the wrong shape, they do not work. Under normal circumstances, the cells will supervise these misfolded proteins and automatically decompose them. However, the supervision mechanism is not with 100% insurance. Scientists found that the rate of decomposition of Prions is not quick enough and these Prions accumulate and change the cellular metabolism and eventually kill the cell. This leads to the death of neurons in the brain. The dead neurons decompose and release more prion proteins into the biological mechanism to cause prion diseases. The infectious diseased prion is thought to be an abnormally folded isoform (PrP^{Sc}) of a host protein known as the prion protein (PrP^C). The conversion of PrP^C to PrP^{Sc} occurs post-translationally and involves conformational change from a predominantly α-helical protein to one rich in β-sheet amyloid fibrils. Much remains to be understood about how the normal cellular isoform of the prion protein PrP^C undergoes structural changes to become the disease associated amyloid fibril form PrP^{Sc}. The "structural conformational" changes of PrP^{Sc} from PrP^C are just very proper to be studied by MD techniques. Classical studies have showed that rabbits have low susceptibility to be infected by PrP^{Sc} [1, 5, 8, 3, 15, 16, 7, 12, 17, 18, 21, 11]. However, recently there is one exception saying that (i) rabbit prion can be produced through saPMCA (*s*erial *a*utomated Protein Misfolding Cyclic Amplification) in vitro [2, 4] (though not by challenging rabbits directly in vivo with other known prion strains), and (ii) the rabbit prion generated is infectious and transmissible [2]. Rabbit prion protein had already had NMR [10] and X-ray [7] structures in Protein Data Bank (www.rcsb.org) with PDB entries 2FJ3, 3O79 respectively. This paper will study the NMR structure of rabbit prion protein (and its I214V and S173N mutants: NMR structure PDB entries 2JOM, 2JOH) by MD techniques, in order to explain the specific mechanism about rabbit PrP^C ($RaPrP^C$) and the conversion of $PrP^C \rightarrow PrP^{Sc}$ of rabbits. Surely, the studies should provide valuable knowledge about the rules governing the $PrP^C \rightarrow PrP^{Sc}$ conversion, which can provide some ideas for designing novel therapeutic approaches that block the conversion and disease propagation.

The rest of this paper is organized as follows. In Section 2, we will briefly review previous research results on rabbit prion protein listed in [1, 5, 8, 3, 15, 16, 7, 12, 17, 18, 21, 11, 2, 4]. Section 3 will present the MD simulation materials and methods for NMR structures of $RaPrP^C$ wild-type and mutants. Section 4 will give MD simulation preliminary results and their preliminary discussions. At last, concluding remarks on RaPrP are summarized in Section 5.

2 Brief Review on Rabbit PrP

The symptoms of TSEs were first described for sheep in 1730 and called “scrapie” in England, “vertige” in France and “Traberkrankheit” in Germany [14]. Now we know that many species such as sheep, goats, mice, humans, chimpanzees, hamsters, cattle, elks, deers, minks, cats, chicken, pigs, turtles, etc are susceptible to TSEs. But many laboratory experiments show that rabbits, horses and dogs seem to be the resistant (or at least the low-susceptibility-rate) species to TSEs. However, recently Chianini *et al.* (2012) reported “rabbits are not resistant to prion infection” [2]. Thus, at this moment it is very necessary for us to briefly review some laboratory works on RaPrP:

- Barlow and Rennie (1976) made many attempts to infect rabbits with the ME7 prion strain and other known prion strains but all failed at last [1].
- Korth *et al.* (1997) found that RaPrP was not recognized by a conformational antibody specific for $\mathrm{PrP^{Sc}}$-like structures [8].
- Vorberg *et al.* (2003) found multiple amino acid residues (such as GLY99, MET108, SER173, ILE214) within the rabbit prion protein inhibit formation of its abnormal isoform [16]. The authors made some substitutions of mouse PrP amino acid sequence by rabbit PrP amino acid sequence and found (i) at the N-terminal region (residues 1–111) the $\mathrm{PrP^{Sc}}$ formation is totally prevented, (ii) at the central region (residues 112–177), the constructed PrP failed to be converted to protease resistance, (iii) at the C-terminal region (residues 178–254) the formation of $\mathrm{PrP^{Sc}}$ is drastically decreased but is not abolished completely. Thus, rabbit cells are negatively affected by the formation of $\mathrm{PrP^{Sc}}$.
- Fernandez-Funez *et al.* (2009) found RaPrP does not induce neurodegeneration in the brains of transgenic flies [5].
- Nisbet *et al.* (2010) created a mutant mouse PrP model containing rabbit PrP-specific amino acids at the GPI anchor site and found that the GPI anchor attachment site (ω site) controls the ability of $\mathrm{PrP^{C}} \rightarrow \mathrm{PrP^{Sc}}$ and the residues at ω and ω+1 of PrP are important modulators of this pathogenic process [12].
- Khan *et al.* (2010) found the propensity to form β-state (the β-sheet-rich structure) is greatest for hamster PrP, less for mouse PrP, but least for the PrP of rabbits, horses and dogs under different conditions and using two-wavelength CD (Circular Dichroism) method they also found a key hydrophobic staple-like helix-capping motif keeping the stability of RaPrP’s X-ray crystallographic molecular structure [7].
- Wen *et al.* (2010) using multidimensional heteronuclear NMR techniques reported that the I214V and S173N substitutions result in distinct structural changes for $\mathrm{RaPrP^{C}}$ [17, 18]. The authors also reported the highly ordered $\beta 2$-$\alpha 2$ loop may contribute to the local as well as global stability of the rabbit PrP protein [18].

- Zhou *et al.* (2011) found that the crowded physiological agents Ficoll 70 and dextran 70 have effects significantly inhibiting fibrillation of rabbit prion protein [21, 11].
- Chianini *et al.* (2012) generated rabbit PrP^{Sc} in vitro subjecting unseeded normal rabbit brain homogenate to serial automated protein misfolding cyclic amplification (saPMCA) and found the rabbit PrP^{Sc} generated in vitro is infectious and transmissible [2]. We noticed that the rabbit prion is just produced through PMCA in vitro not by challenging rabbits directly in vivo with other known prion strains.

3 Materials and Methods

This is the continuation and extension of the MD simulations of [19]. Zhang (2010) carried out (i) 15 ns of production phase of MD simulations and (ii) the heating phase of MD simulations is starting from 100 K with one set of initial velocity (denoted as seed2) [19]. This paper continued to finish another 15 ns of production phase of MD simulations for seed2, and carried out other two sets of initial velocities for heating from 100 K (denoted as seed1 and seed3) of MD simulations with 30 ns of production phases.

4 Preliminary Results and Preliminary Discussions

For seed1$\sim$seed3, during the whole 30 ns, radii of gyrations of the wild-type and the I214V and S173N mutants have been always level off around 15 angstroms whether in neutral pH environment or in low pH environment.

In neutral pH environment, compared with the I214V and S173N mutants, the wild-type is always at the lowest level of RMSD (root mean square deviation), RMSF (root mean square fluctuation) and B-factor values. This show to us the wild-type RaPrP is very stable under neutral pH environment. However, under low pH environment, we cannot see some large differences among the wild-type, the I214V mutant and the S173N mutant.

The variations of the molecular structures of the wild-type and its I214V and S173N mutants during the 30 ns for the three sets of MD simulations can be seen in Figures 1$\sim$3, where the graphs were produced in the use of DSSP program [6], H is the α-helix, I is the π-helix, G is the 3-helix or 3/10 helix, B is the residue in isolated β-bridge, E is the extended strand (participates in β-ladder), T is the hydrogen bonded turn, and S is the bend; H1, H2, and H3 respectively denote the α-helix 1, α-helix 2 and α-helix 3 of a prion protein. We can see that for the wild-type the three α-helices have been still kept during the whole 30 ns of each set under the neutral pH environment, but under low pH environment, the three α-helices of the wild-type are unfolded as of the I214V and S173N mutants under neutral or low pH environment and at last should be unfolded into rich β-sheet structures.

Under low pH environment, the salt bridges of the wild-type (Table 1) are broken thus leads to the unfolding of the stable helical structures of RaPrP. The following salt bridges should contribute to the stability of RaPrP: ASP146-ARG147, GLU210-ARG207, GLU206-LYS203, GLU206-ARG207, ASP146-HIS139, GLU151-ARG150, GLU151-ARG147, GLU151-ARG155, ASP177-ARG163, GLU145-ARG135, ASP143-HIS139, GLU145-HIS139, GLU195-ARG155, ASP146-ARG150, ASP177-HIS176, ASP143-ARG147, GLU195-LYS193, HIS186-ARG155, HIS186-LYS184, ASP166-ARG163, GLU210-HIS176, HIS139-ARG135, GLU199-LYS193, GLU206-HIS176. Among these salt bridges, compared with the salt bridges of I214V mutant and S173N mutant listed in Table 1 and the observation from Figures 1$\sim$3, we find that GLU-195-ARG155 and ASP166-ARG163 are important to contribute to the structural stability of wild-type RaPrP.

Zhang (2011) found that there always exist salt bridges between ASP202-ARG156, ASP178-ARG164 in human and mouse prion proteins, between ASP201-ARG155, ASP177-ARG163 in RaPrP [20]. The author broke salt bridges of of human, mouse and rabbit prion proteins by doing MD simulations from neutral to low pH environment; consequently, the secondary structures of human and mouse prion proteins were not changed very much, while the stable helical structure of wild-type RaPrP collapsed [20]. This is to say that the human and mouse prion protein structures are not affected by removing these salt bridges but the structure of RaPrP is affected very much by these salt bridges. Therefore, compared with human and mouse prion proteins, the salt bridges such as ASP201-ARG155, ASP177-ARG163 contribute greatly to the structural stability of RaPrP.

5 A Concluding Remark

Like the controversy on “prion” theory, there is also a big controversy over “whether rabbits are resistant to prion infection or not?”. This paper briefly reviewed the research results on rabbit prion protein and rabbit prions. This paper also did MD simulation studies on RaPrP and its I214V and S173N mutants with a little preliminary analysis of the role of salt bridges that plays in the structural stability of RaPrP. Further studies to understand the specific mechanism of RaPrP are clearly needed.

Acknowledgments

This research has been supported by a Victorian Life Sciences Computation Initiative (VLSCI) grant numbered VR0063 on its Peak Computing Facility at the University of Melbourne, an initiative of the Victorian Government of Australia.

References

[1] Barlow RM, Rennie JC (1976) The fate of ME7 scrapie infection in rats, guinea-pigs and rabbits. Res Vet Sci 21: 110–111.

[2] Chianini F, Fernndez-Borges N, Vidal E, Gibbard L, Pintado B, de Castro J, Priola SA, Hamilton S, Eatona LS, Finlayson J, Pang Y, Steele P, Reid HW, Dagleish MP, Castilla J (2012) Rabbits are not resistant to prion infection. Proc Natl Acad Sci USA 109(13): 5080-5085.

[3] Courageot MP, Daude N, Nonno R, Paquet S, Di Bari MA, Le Dur A, Chapuis J, Hill AF, Agrimi U, Laude H, Vilette D (2008) A cell line infectible by prion strains from different species. J Gen Virol 89: 341–347.

[4] Fernandez-Borges N, Chianini F, Erana H, Vidal E, Eaton SL, Pintado B, Finlayson J, Dagleish MP, Castilla J (2012) Naturally prion resistant mammals: A utopia? Prion 6(5): 425–429.

[5] Fernandez-Funez P, Casas-Tinto S, Zhang Y, Gomez-Velazquez M, Morales-Garza MA, Cepeda-Nieto AC, Castilla J, Soto C, Rincon-Limas DE (2009) In vivo generation of neurotoxic prion protein: role for hsp70 in accumulation of misfolded isoforms. PLoS Genet 5: e1000507.

[6] Kabsch W, Sander C (1983) Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. Biopolymers 22: 2577-2637.

[7] Khan MQ, Sweeting B, Mulligan VK, Arslan PE, Cashman NR, Pai EF, Chakrabartty A (2010) Prion disease susceptibility is affected by β-structure folding propensity and local side-chain interactions in PrP. Proc Natl Acad Sci USA 107: 19808-19813.

[8] Korth C, Stierli B, Streit P, Moser M, Schaller O, Fischer R, Schulz-Schaeffer W, Kretzschmar H, Raeber A, Braun U, Ehrensperger F, Hornemann S, Glockshuber R, Riek R, Billeter M, Wthrich K, Oesch B (1997) Prion ($\mathrm{PrP^{Sc}}$)-specific epitope defined by a monoclonal antibody. Nature 390: 74–77.

[9] Lin DH, Wen Y (2011) Progresses on prion proteins. Scientia Sinica Chimica 41(4): 683–698.

[10] Li J, Mei FH, Xiao GF, Guo CY, Lin DH (2007) 1^{H}, 13^{C} and 15^{N} resonance assignments of rabbit prion protein 91–228. J Biol NMR 38: 181.

[11] Ma Q, Fan JB, Zhou Z, Zhou BR, Meng SR, Hu JY, Chen J, Liang Y (2012) Fibril Formation of the Rabbit/Human/Bovine Prion Proteins. Biophys J 101: 1483–1492.

[12] Nisbet RM, Harrison CF, Lawson VA, Masters CL, Cappai R, Hill AF (2010) Residues surrounding the glycosylphosphatidylinositol anchor attachment site of PrP modulate prion infection: insight from the resistance of rabbits to prion disease. J Virol 84(13): 6678–6686.

[13] Polymenidou M, Trusheim H, Stallmach L, Moos R, Julius C, Miele G, Lenz-Bauer C, Aguzzi A (2008) Canine MDCK cell lines are refractory toinfection with human and mouse prions. Vaccine 26(21): 2601-2614.

[14] Verdier J-M (2012) Prions and Prion Diseases: New Developments, NOVA Science Publishers, ISBN 978-1-62100-027-3, Hard cover, Preface, pp. vii-ix.

[15] Vilette D, Andreoletti O, Archer F, Madelaine MF, Vilotte JL, Lehmann S, Laude H (2001) Ex vivo propagation of infectious sheep scrapie agent in heterologous epithelial cells expressing ovine prion protein. Proc Natl Acad Sci USA 98: 4055–4059.

[16] Vorberg I, Martin HG, Eberhard P, Suzette AP (2003) Multiple amino acid residues within the rabbit prion protein inhibit formation of its abnormal isoform. J Virol 77: 2003–2009.

[17] Wen Y, Li J, Xiong M, Peng Y, Yao W, Hong J, Lin DH (2010) Solution structure and dynamics of the I214V mutant of the rabbit prion protein. PLoS ONE 5: e13273.

[18] Wen Y, Li J, Yao W, Xiong M, Hong J, Peng Y, Xiao G, Lin DH (2010) Unique structural characteristics of the rabbit prion protein. J Biol Chem 285: 31682–31693.

[19] Zhang JP (2010) Studies on the structural stability of rabbit prion protein probed by molecular dynamics simulations of its wild-type and mutants. J Theor Biol 264: 119–122.

[20] Zhang JP (2011) Comparison studies of the structural stability of rabbit prion protein with human and mouse prion proteins. J Theor Biol 269: 88–95.

[21] Zhou Z, Yan X, Pan K, Chen J, Xie ZS, Xiao GF, Yang FQ, Liang Y (2011) Fibril Formation of the Rabbit/Human/Bovine Prion Proteins. Biophys J 101: 1483–1492.



Figure 1: *(Seed1) Secondary Structure graphs for rabbit prion protein wild-type, I214V mutant and S173N mutant (from up to down) (x-axis: time (0–30 ns), y-axis: residue number (124–228); left column: neutral pH, right column: low pH.*



Figure 2: *(Seed2) Secondary Structure graphs for rabbit prion protein wild-type, I214V mutant and S173N mutant (from up to down) (x-axis: time (0–30 ns), y-axis: residue number (124–228); left column: neutral pH, right column: low pH.*



Figure 3: *(Seed3) Secondary Structure graphs for rabbit prion protein wild-type, I214V mutant and S173N mutant (from up to down) (x-axis: time (0–30 ns), y-axis: residue number (124–228); left column: neutral pH, right column: low pH.*

Table 1: Salt bridges of the wild-type and the I214V and S173N mutants under neutral pH environment for the MD simulations with occupied rates for seed1, seed2, seed3, where * denotes the occupied rate is less than 10%.

Salt bridges \	wild-type	I214V mutant	S173N mutant
ASP146-ARG147	(100, 100, 100)	(100, 100, 99.90)	(100, 99.97, 100)
GLU210-ARG207	(99.15, 97.58, 98.43)	(35.08, 98.08, *)	(99.25, 55.00, 100)
GLU206-LYS203	(97.67, 92.85, 87.35)	(29.07, 41.00, 57.78)	(*, 32.52, 46.10)
GLU206-ARG207	(84.25, 84.80, 75.50)	(65.78, 57.03, 73.68)	(47.33, 27.07, 18.75)
ASP146-HIS139	(*, 73.65, *)	(69.13, *, *)	(*, 38.93, *)
GLU151-ARG150	(72.13, 47.10, 84.25)	(65.35, 68.42, 88.25)	(57.32, 67.85, 75.15)
GLU151-ARG147	(63.13, 36.28, 64.22)	(39.02, 37.32, 34.45)	(*,18.67, 73.33)
GLU151-ARG155	(60.78, *, 60.07)		(*, *, 29.00)
ASP177-ARG163	(47.80, 40.38, 21.92)	(*,25.17, *)	
GLU145-ARG135	(33.52, *, 47.13)	(*, *, 13.30)	
ASP143-HIS139	(27.95, *, 10.95)	(*, *, 29.40)	(*, 26.97, 13.53)
GLU145-HIS139	(25.00, *, 18.60)		(*, *, 26.72)
GLU195-ARG155	(21.88, *, *)		
ASP146-ARG150	(21.23, 66.38, 34.25)	(59.57, 15.38, 29.25)	(42.77, 66.85, 12.42)
ASP177-HIS176	(21.17, 16.55, 10.15)	(70.05, 42.13, 20.30)	(25.53, 34.40, *)
ASP143-ARG147	(19.73, 50.30, 28.38)	(33.85, *, 21.23)	(10.43, 42.20, 61.53)
GLU195-LYS193	(19.68, 21.98, 10.27)	(*, 19.53, *)	(26.15, 20.60, *)
HIS186-ARG155	(14.01, 96.35, 12.47)	(*, *, 11.37)	
HIS186-LYS184	(*, 37.95, 12.95)	(*, 30.79, *)	(82.28, 86.89, *)
ASP166-ARG163	(*, *, 54.08)		
GLU210-HIS176	(*, *, 33.37)		(24.25, *, *)
HIS139-ARG135	(*, *, 32.02)		(*, *, 54.32)
GLU199-LYS193	(*, *, 11.95)	(17.77, *, *)	(*, 18.33, *)
GLU206-HIS176	(*, *, 11.83)	(*, 62.37, *)	
HIS176-ARG163		(72.11, *, *)	(*, 39.57, *)
ASP201-LYS203		(32.53, 87.10, 91.17)	(38.32, 69.48, 73.22)
GLU199-LYS184		(25.53, *, *)	(18.22, *, *)
GLU195-HIS186		(23.32, *, *)	(*, 25.82, 16.67)
HIS186-LYS193		(27.65, *, *)	(*, 10.70, *)
GLU145-ARG147		(*, 80.58, 24.40)	
GLU210-HIS139		(*, 28.42, *)	
ASP201-ARG155		(*, 27.60, *)	
HIS176-HIS186		(*, 20.18, *)	
GLU220-ARG227		(*, 10.60, *)	
GLU206-LYS184		(*, *, 46.75)	
ASP177-ARG207		(*, *, 40.75)	
HIS176-ARG207		(*, *, 15.67)	
GLU151-LYS193		(*, *, 14.55)	
GLU206-HIS186		(*, *, 13.08)	
ASP143-ARG135			(24.60, *, *)
GLU199-HIS186			(18.68, *, *)
GLU206-ARG155			(17.50, *, *)
ASP201-HIS186			(14.48, *, *)
GLU145-ARG150			(13.33, *, *)
HIS176-LYS184			(*, 21.93, *)
GLU195-ARG227			(*, 12.28, *)
ASP146-ARG135			(*, 10.70, *)
HIS176-ARG227			(*, *, 11.12)
GLU220-HIS176			(*, *, 24.17)
ASP201-ARG207			(*, *, 15.55)
GLU210-ARG227			(*, *, 12.00)